

4th International Workshop on the CKM Unitarity Triangle (CKM2006)

Proposed Workshop Format

Note:

- Convener candidates are listed in order of priority for each exp. and theory.
- Conveners at CKM2005 at UCSD are marked by *.

WG1: Precise Determination of V_{ud} and V_{us}

Conveners: 2 (exp.) + 2 (theory)

IOC Contacts: S.Hashimoto, T.Iijima, T.Komatsubara, R.Briere

The main topic is the precise determination of the Cabibbo angle through the determination of the CKM elements V_{ud} and V_{us}

- Determination of V_{us}
 - from K_{13} decays: new K_{13} results, measurement and analysis of K_{13} Dalitz plot to constrain CHPT parameters, new lattice results and their errors
 - from τ and hyperon decays: estimate of theoretical errors, prospects
 - from f_K/f_π : new lattice results, exhaustive estimate of unquenched lattice uncertainty
- Determination of V_{ud} from super-allowed Fermi transitions, neutron and pion decays: theoretical and experimental updates, also new idea.
- Determination of V_{cs} , V_{cd}

WG2: Determination of V_{ub} and V_{cb} through Inclusive / Exclusive Semileptonic B decays

Conveners: 2(exp.)+2(theory)

IOC Contacts: P.Gambino, S.Hashimoto, A.Stocchi, R.Briere

The main goal of this group is the determination of V_{ub} and V_{cb} through the inclusive and exclusive semileptonic B decays

- Determination of V_{cb} and HQE parameters from inclusive $B \rightarrow X l \nu$ decay.
 - Theoretical uncertainties: prospects
 - Spectral moments: new results, averages, global fits, scheme differences, determination of quark masses and hadronic parameters and their combination.
- Determination of V_{ub} inclusive
 - New measurements of E_{lepton} , M_X , q^2 spectrum.
 - Methods to determine $|V_{ub}|$ from the measured rate, based on shape function scheme, weighting functions and DGE(Dressed Gluon Exponentiation) etc.
 - Associated theoretical errors. Weak annihilation effect.
 - How to make the average etc.
- Determination of V_{cb} exclusive: theoretical and experimental update
 - New measurements for the form factors.
 - $F(1)$ lattice unquenched calculation, LCSR
- Determination of V_{ub} exclusive: theoretical and experimental update
 - New measurements of the q^2 distribution with different tagging methods.
 - Experimental determination of the form factor shape.
 - Form factor normalization from unquenched lattice, LCSR and others.
- Exclusive D semileptonic decays and form factors
- $D \rightarrow \pi l \nu$, extrapolation from D to B and possible determination of V_{ub}

WG3: V_{td}/V_{ts} through Mixings and Rare B and K Decays

Conveners: 2(exp.)+2(theory)

IOC Contacts: P.Gambino, T.Komatsubara, F.Ukegawa, R.Forty

The main goal is determination of V_{td}/V_{ts} through meson mixing and rare B and K decays.

- ϵ_K , ΔM_d , ΔM_s .
 - Lattice determination and experimental constraints on F_{Bd} , F_{Bs} , and ζ
 - Critical review on quenching and chiral logs associated errors
- B hadron widths and differences: theoretical and experimental update
- Measurement of $D_s/D^+ \rightarrow \tau(\mu)\nu$

- Determination of f_{D_s} , f_{D^+} and their ratio
- Possible extrapolation to f_B and ζ
- V_{td}/V_{ts} from $B \rightarrow \rho\gamma$ and $B \rightarrow K^*\gamma$
 - New results from experiments.
 - Results from lattice, LCSR, prospects, long distance contributions
- Radiative decays as probe of CKM
 - Theoretical and experimental updates of $B \rightarrow X_s\gamma$
 - use of $b \rightarrow s\gamma$ for moments and for V_{ub} (discussion in conjunction with WG2)
- Role of the radiative charm decays.
- Rare K decays of relevance to CKM studies

WG4: Angles from charmless B decays

Conveners: 2(exp.) + 2(theory)

IOC Contacts: R.Faccini, M.Hazumi, T.Iijima, V.Sharma

The main goal is to study the measurements of the angles ϕ_1, ϕ_2, ϕ_3 (or β, α, γ) in charmless B decays. The limitations and ways to overcome them will be discussed, as well as new approaches and high statistics projections. We will concentrate on ways to estimate the SM uncertainties in all these modes and theoretical and experimental ways to reduce them.

- ϕ_2 (α) modes: two body decays ($\pi\pi$), quasi-two-body (Q2B) B decays ($\rho\rho, \rho\pi$) and Dalitz analyses ($\pi\pi\pi$).
 - New results ? New modes ?
 - How can we get to $f_2(\alpha)$ from the different measurements? How good is the Q2B approximation? How important are S-wave contribution (and other interference) in modes involving ρ ?
 - How large are isospin breaking effects? Are there ways that these errors can be reduced?
 - How trustable are theoretical calculations of the penguin pollution?
- ϕ_1 (β) modes: $\phi K_0, \eta' K_0, f_0 K_0, \pi_0 K_S, K+K-K_S, K_S K_S K_0$ and more.
 - New results ? New modes ?
 - Possible deviation from $\sin 2\phi_1$ ($\sin 2\beta$) within the SM? What are the methods one can use to calculate them and how these methods can be checked?

- Can the sign of the deviation be obtained ?
- Are there correlations between the deviation in these modes? Are there observables or combinations of observables that have smaller SM uncertainties (sums, ratios etc.) ?
- ϕ_3 (γ) from charmless decays. Interpretation of other charmless decays
 - Understanding of the πK modes (γ , electroweak penguins).
 - What can be learnt in addition from πK^* and ρK^* modes?
- Comparison and assessment of theoretical approaches based on factorization and/or flavour symmetries
 - What are the theoretical assumptions?
 - Which amplitudes are included in the calculations, and what is the estimated theoretical error?
 - Perform comparative studies for “benchmark” modes.
 - How predictive are data-driven fits?

WG5: Angles from B decays with charm

Conveners: 2(exp.) + 2(theory)

IOC Contacts: R.Faccini, M.Hazumi, Y.Sakai, V.Sharma, A.Stocchi

The main goal is the measurement of the angles ϕ_3 and β from B decays involving D or Charmonium mesons. The limitations and ways to overcome them will be discussed, as well as new approaches and high statistics projections

- Measurement of ϕ_1 (β) in B decays to Charmonium
 - measurements of $\phi_1(\beta)$ and $\cos 2\phi_1$ ($\cos 2\beta$)
 - CPT/T parameters
- Measurements of ϕ_3 (γ) in $B \rightarrow D^{(*)}K^{(*)}$ decays
 - GLW and ADS methods
 - Dalitz methods (model dependent/independent)
- Measurements of $2\phi_1 + \phi_3$ ($2\beta + \gamma$) in $B \rightarrow D^{(*)}\pi, \rho$ decays and related BF measurements ($D_s\pi, D\pi^0$, and $D^{(*)}K^{(*)}$)

WG6: CKM Fits and New Physics

Conveners: 2(exp.) + 2(theory)

IOC Contacts: Y.Sakai, A.Stocchi, F.Ukegawa, R.Forty

The main goals are global fits to the CKM parameters using different statistical methods, and search for, or constrain, the new physics contribution, using data from the B factories, Tevatron, and the future B facilities.

- Global fit to Unitarity Triangle parameters using all the experimental and theoretical inputs from all the other WGs
 - Comparison between direct and indirect determinations of different quantities, for instance, UT angles, hadronic parameters.
 - $f_B^* |V_{ub}|$ from $B \rightarrow \tau \nu$, $|V_{ub}/V_{td}|$ from $B \rightarrow \tau \nu$ and Δm .
- Constraint to New Physics from
 - FCNC processes such as $B \rightarrow X_s \gamma$, X_{sll} , $X_{s\nu\nu}$, $K \rightarrow \pi \nu \nu$ etc.
 - B decays to a final state with τ , such as $B \rightarrow D \tau \nu$, $\tau \nu$.
 - Decays via Higgs mediation, such as $B_s \rightarrow \mu^+ \mu^-$.
 - Others.